

A SACROILIAC JOINT INJECTION *is an outpatient procedure for treating low back and buttock pain.*

This information has been provided by your doctor so you can better understand this procedure. Your doctor will make the best recommendation for your specific needs.

This pamphlet is for general education only. Specific questions or concerns should always be directed to your doctor. Your doctor can explain possible risks or side effects.



(C) Copyright OEA For OEA website use only

This region is blank on the printed brochures allowing you to place your business card label here. This area can also be customized with your clinic information. See the customized brochures tab on the OEA website: www.oeabrochures.com

SACROILIAC JOINT INJECTION

for Chronic Low Back and Buttock Pain



An outpatient procedure for treating low back and buttock pain




OEA PAIN BROCHURES
www.oeabrochures.com

Copyright © 2005-2009 OEA, LLC. INV-0709

Sacroiliac Joint Injection for Chronic Low Back and Buttock Pain

WHAT ARE SACROILIAC JOINTS?

Sacroiliac joints connect your spine to your hip bone. They connect the bottom of the spine, called the sacrum, to the outer part of the hip bone, called the ilium. You have two sacroiliac joints. One is found on each side of the sacrum.



Sacroiliac joints help control your pelvis when you move. They help transfer forces from your lower body to your upper body. Each sacroiliac joint has several ligaments to help strengthen it.

WHAT IS SACROILIAC JOINT PAIN?

You may feel pain if a sacroiliac joint is injured. At times it may feel like simple muscle tension, but at other times the pain can be severe.

Sometimes the cartilage inside the joint is injured, but at other times only the ligaments around the joint are affected.


You usually feel sacroiliac joint pain in an area from your low back down to your buttocks. But sometimes, if a joint is very inflamed, pain may even extend down the back of the leg. The diagram shows where sacroiliac joint pain usually is felt.

HOW DO I KNOW IF I HAVE SACROILIAC JOINT PAIN?


If you have pain in one or more of these areas you may have sacroiliac joint pain. Common tests such as x-rays or MRIs may not always show if a sacroiliac joint is causing pain. Your pain doctor can perform other tests to find out if you have sacroiliac joint pain.

WHAT IS A SACROILIAC JOINT INJECTION?

In a sacroiliac joint injection, a local anesthetic and corticosteroid are injected into one or both of your sacroiliac joints, or into the ligaments surrounding the joints. The local anesthetic, a numbing medicine, lessens your pain temporarily. The corticosteroid reduces inflammation that may be causing pain.



The injection can be used for diagnosis and treatment. If the injection immediately lessens your pain and helps you move better, it tells the doctor which joint is causing the pain.



Fortin JD, Dwyer AP, West S, Pier J. Sacroiliac Joint Pain Referral Maps Upon Applying a new Injection/Arthrography. Spine. 1994; 19:1475-1482

WHAT HAPPENS AFTER AN INJECTION?

You will be monitored for up to 30 minutes after the injection. When you are ready to leave, the clinic will give you discharge instructions. You will also be given a pain diary. It is important to fill this out because it helps your doctor know how the injection is working.

It may help to move in ways that hurt before the injection, to see if the pain is still there, but do not overdo it. Take it easy for the rest of the day.

You may feel immediate pain relief and numbness in your back for up to six hours after the injection. This tells you the medication has reached the right spot.

Your pain may return after this short pain-free period, or may even be a little worse for a day or two. This is normal. It may be caused by needle irritation or by the corticosteroid itself. Corticosteroids usually take two or three days to start working, but can take as long as a week.

You can usually return to work the day after the injection, but always check with your doctor.

HOW LONG CAN I EXPECT PAIN RELIEF?

How long you can expect pain relief depends on how many areas are injured and on the amount of inflammation.

Sometimes an injection can bring several months of pain relief, and then more treatment is needed. Other times, particularly if there is no underlying bone or joint problem, one injection brings long-term pain relief. If your pain is caused by injury to more than one area, only some of your symptoms may be helped by one injection.

